

Supplementary Table 8. Characteristics of included studies in key question #7

Studies	Patients	Arm A	Arm B	Arm-C	Duration	Outcome 1
Hawkey et al. ⁶⁶ (1998)	Taking NSAIDs and ulcers or more than 10 erosions	Omeprazole 20 mg/D (n=308)	Omeprazole 40 mg/D (n=315)	Misoprostol 200 ug qid (n=298)	If endoscopic gross erosions remain after 4 weeks, repeat EGD after 8 weeks and maintenance for 6 months	Endoscopic healing
Yeomans et al. ⁶⁷ (1998)	Taking NSAIDs and ulcers or more than 10 erosions	Omeprazole 20 mg/D (n=174)	Omeprazole 40 mg/D (n=187)	Ranitidine 150 mg bid (n=174)	If endoscopic gross erosions remain after 4 weeks, repeat EGD after 8 weeks and maintenance for 6 months	Endoscopic healing
Karateev and Murav'ev ⁶⁸ (2001)	Taking NSAIDs and multiple erosion, ulcer, or duodenal ulcer	X	Omeprazole 40 mg/D (n=36)	Misoprostol 200 ug qid (n=27)	2 weeks	Endoscopic healing

NSAID, non-steroidal anti-inflammatory drug; qid, four times a day; bid, twice a day; EGD, esophagogastroduodenoscopy.